

CLINICAL PRACTICE GUIDELINES

JUNE 2013

MOH/P/PAK/266.13[GU]

MANAGEMENT OF PSORIASIS VULGARIS



Ministry of Health Malaysia



Dermatological Society of Malaysia



Academy of Medicine Malaysia

Systemic Therapy

Clinical Practice Guidelines Management of Psoriasis Development Group

Dr. Ambrose
Department of Medicine
Hospital Putrajaya

Systemic Therapy

- Moderate to severe psoriasis frequently require **systemic or biologic therapy**.

- Systemic agents :

- **methotrexate**
- **acitretin**
- **cyclosporine**

have significant side-effects and cumulative toxicity

- **Pretreatment** assessment

- to identify risk of developing toxicity.

- Baseline laboratory/ imaging tests

- regularly to monitor side effects/ toxicity

Malaysian CPG on the management of Psoriasis vulgaris

Pre-treatment assessment

RECOMMENDATION

- All patient for systemic/biologic therapy should have a pre-treatment assessment including laboratory / imaging tests and regular monitoring for side effects/toxicity*. **(Grade C)**

Methotrexate

- Methotrexate
 - analogue of folic acid
 - inhibits dihydrofolate reductase
- Methotrexate is efficacious in treating moderate to severe plaque psoriasis
- Bansback et al meta analysis on efficacy of systemic therapy
 - 42% of patients on methotrexate 15 to 22.5 mg/week
 - PASI 75 at week 16
 - RR of 9.8 (95% CI 6.08 to 13.19)
 - NNT=3
- In a separate study
 - 70% of patients taking methotrexate 15 to 20mg/week
 - PASI 75 at week 12
- No benefit in increasing methotrexate from 20 to 25 mg/week in patients who failed to achieve PASI 50 at week 12

Methotrexate

- No significant difference in patients taking methotrexate 15 to 22.5 mg/week compared to cyclosporine (60.5% vs 71.4%) ($p=0.09$)
- Methotrexate was comparable to mycophenolate mofetil with 73.3% vs. 58.8% of patients achieving PASI 75 at week 12 ($p>0.05$)
- Methotrexate was superior to hydroxycarbamid

Methotrexate

- Adverse events:
 - hepatotoxicity
 - myelosuppression
 - gastrointestinal manifestation
 - nausea
 - vomiting
 - mouth sores
 - loss of appetite
 - hair loss and malaise
- Most serious side effects are
 - hepatotoxicity
 - myelosuppression
- Identify the risk factors before initiation of methotrexate

Methotrexate

- Severity of side-effects is dose dependant
- Most common cause of withdrawal : hepatic adverse events
- Common gastrointestinal side effects
 - were nausea
 - diarrhoea
 - abdominal pain
- Incidence of liver fibrosis ranged from 5.7% to 71.8%
- risk factors for liver fibrosis:
 - type 2 diabetes mellitus (OR=7.7, 95%CI 2.7 to 21.7)
 - obesity (OR=2.4, 95% CI 1.1 to 5.4)
- Alcohol consumption and viral hepatitis were **not** significant risk factors

Methotrexate

- Data on myelosuppression with methotrexate were derived from patients with rheumatoid arthritis
- The risk of myelosuppression in psoriasis patients is unknown

Risk factors for methotrexate induced hematotoxicity

- Renal insufficiency
- Advanced age
- Lack of folate supplementation
- Medication errors
- Drug interactions
- Hypoalbuminemia
- Excess alcohol intake

Methotrexate

- Monitoring of hepatotoxicity varies in different centres.
 - regular serum liver function test to liver biopsy.
- Routine to do liver biopsy after a cumulative methotrexate dose of 1.5 gm ---→1.0 to 1.5 gm interval to monitor methotrexate-induced hepatotoxicity.
- Recent data : risk of developing liver fibrosis is less than 2.6% in **low risk patients** taking a cumulative methotrexate dose of more than 4g.
- Liver biopsy deferred till a cumulative dose of ≥ 4 gm
 - in low risk patients taking low dose (<20 mg) methotrexate.

Methotrexate

- Non-invasive methods to monitor hepatotoxicity such as
 - Serum Procollagen III aminopeptide
 - (sensitivity : 77.3%, specificity : 91.5%)
 - fibrotest
 - (sensitivity : 83% , specificity : 61 %)
 - Fibroscan
 - (sensitivity : 50%, specificity : 88%)

NOT widely available in Malaysia.

- A liver biopsy can be deferred if the level of **procollagen type III** remains within the normal limits

Methotrexate

- Supplementation with folic acid or folinic acid is an effective measure to reduce hepatic adverse effects
- No significant reduction in side effects of
 - Gastrointestinal (ARR= -0.09, 95% CI -0.2 to 0.02)
 - Mucocutaneous (ARR= -0.07, 95% CI -0.2 to 0.04)
 - Haematological (ARR=0.004, 95% CI -0.02 to 0.03)

Methotrexate

- Data on risk of pulmonary fibrosis in long term MTX is limited
- A study on 27 psoriatic arthritis patients on low dose MTX (5 - 15 mg/week for 52 months showed no association with pulmonary fibrosis
- Systematic review reported 84 cases of lung related adverse event in 3463 patients with rheumatoid arthritis on methotrexate
 - Only 15 of which were felt to be directly caused by methotrexate (incidence 0.43%).

Treatment regime and monitoring of Methotrexate in patients with psoriasis

Initial therapy


- oral test dose of 5.0 - 7.5 mg /week
- Supplement with Folic acid 5mg od (except the day of methotrexate) or 5mg once a week (the day after methotrexate)
- Repeat FBC, LFT and RP within 2 weeks

Maintenance therapy

- Escalate dose from 7.5 mg/week till response (maximum 20 mg/week)
[administered as a single dose or divided into 3 doses and administered at 12-h intervals over 2 consecutive days]
- Monitor FBC/LFT/RP monthly for the first 3 months and then after every 1 to 3 months
- More frequent monitoring is needed with dose escalation
- Do blood test 5 - 7 days after last dose of methotrexate
- Monitor total cumulative dose of methotrexate
 - Consider Procollagen III aminopeptide/ Fibroscan/Fibrotest or Liver biopsy when achieve total cumulative dose of 3.5 to 4.0g (without risk factors for hepatotoxicity) or 1.0 to 1.5g (with risk factors for hepatotoxicity)*

*Refer to yellow box on “Risk factors for methotrexate induced hepatotoxicity”

ALGORITHM 3: MONITORING HEMATOTOXICITY AND HEPATOTOXICITY OF METHOTREXATE



Malaysian CPG on the management of Psoriasis vulgaris **Methotrexate**

RECOMMENDATION

- Methotrexate should be used as first-line systemic treatment for moderate to severe plaque psoriasis. **(Grade A)**
- Neutropaenia and hepatotoxicity should be monitored closely. **(Grade C)**

Retinoids (Acitretin)

- Retinoids are vitamin A analogues
 - modulate epidermal proliferation and differentiation.
- First study to be published on acitretin was in 1984
- No recent RCTs after 2000 in assessing efficacy of oral retinoids

Retinoids (Acitretin)

- A study on adverse effects of acitretin by Pearce et al. Showed
 - low dose acitretin (<25 mg/day) had less adverse effects
 - compared to high dose acitretin (>25 mg/day)
- The side-effects were
 - cheilitis
 - skin peeling
 - pruritus, alopecia
 - dry mouth
 - xerophthalmia
 - raised alanine transaminase, aspartate aminotransferase and triglycerides
- When phototherapy is not an option, Acitretin is an appropriate alternative treatment for patients with psoriasis

Retinoids (Acitretin)

Initial therapy

- Start with dose of 0.5 – 1 mg/kg/day for 2 - 4 weeks
- Baseline lipid profile and LFTs

Maintenance therapy

- Adjust according to response, usually within range of 25 - 50 mg daily (max 75 mg daily).
- Repeat lipid profile and LFTs every 4 - 8 weeks until stable, then every 6-12 weeks

**in rare cases of use in women of childbearing age, a baseline pregnancy test should be done and repeated monthly*

Malaysian CPG on the management of Psoriasis vulgaris

Acitretin

RECOMMENDATION

- Acitretin may be offered for the treatment of moderate to severe plaque psoriasis. **(Grade A)**
- Acitretin should be avoided in women of childbearing age without reliable contraception and in those who are planning pregnancy however; it is safe for men who are planning to have a child. **(Grade C)**

Cyclosporine

- Cyclosporine is an oral calcineurin inhibitor with cyclical peptide composed of 11 aminoacids and strongly hydrophobic
- Cyclosporine is efficacious as shown by Schmitt et al
 - PASI 75, at week 8 to 16 ranged from 28% to 97%
 - RD of 0.3 (95% CI 0.1 to 0.5)
- This wide range of efficacy was due to different initial dosage of cyclosporine

Cyclosporine

- Adverse events of 16.1%
- Smaller percentages with serious adverse events (2.3%)
- Withdrawal due to adverse events (1.2%)
- In a 5-year cohort study on the risk of malignancy
 - risk of non-melanoma skin malignancies was higher among patients treated for more than 2 years
(SIR >2 years=11.4, 95% CI 5.2 to 21.7 vs
SIR <2 years= 4.6, 95% CI 2.4 to 8.1)
- Previous exposure to PUVA increased the risk of non-melanoma skin cancer
 - (RR = 7.3, 95% CI 1.3 to 134.5)

Cyclosporine

Treatment regime and monitoring of cyclosporine in patients with psoriasis

Pre Cyclosporine screening:

- **Ensure normal baseline Investigations (refer appendix 7) prior to cyclosporine**
- **Discuss benefit & risk of cyclosporine with patient**

Initial therapy

- **Starting dose of 2.5 mg/kg/d divided twice a day**
- **If inadequate response to initial dose after 4 to 6 weeks, the dose may be increased to a maximum of 6 mg/kg of body weight.**
- **Treatment for more than 2 years not recommended**
- **Monitoring while on therapy:**
 - **Blood pressure, RP, FBC, uric acid, potassium, lipids, liver enzymes, serum bilirubin and magnesium should be monitored monthly**

Malaysian CPG on the management of Psoriasis vulgaris Cyclosporine

RECOMMENDATION

- Cyclosporine may be offered as short-term treatment for rapid disease clearance in moderate to severe psoriasis. **(Grade A)**
- Cyclosporine may be offered as second-line systemic agent to psoriasis patients who fail, are intolerant or have contraindications to methotrexate. **(Grade A)**
- Cyclosporine should **NOT** be used for more than 2 years. **(Grade B)**
- Cyclosporine should be avoided in patient with previous PUVA exposure. **(Grade B)**
- Blood pressure, renal function, lipid profile should be monitored closely in psoriasis patient on cyclosporine. **(Grade C)**

Mycophenolate Mofetil

- Mycophenolate mofetil (MMF) is an immune modulator which inhibits inositol monophosphate dehydrogenase.
- MMF (2 gm daily) is **less efficacious** than cyclosporine (2.5 mg/kg/day) with PASI 75 at week 12 of 12% and 29% respectively ($p=0.01$)
- Mean PASI improvement is lower in MMF (30 mg/kg/day) compared to cyclosporine (4 mg/kg/day) ($p=0.04$)
- The tolerability and adverse events are similar between the two drugs

Antibiotics

- There is no good evidence to support the use of antibiotics in treating plaque psoriasis

Hydroxyurea

- Hydroxyurea is an antimetabolite agent which inhibits deoxyribonucleic acid.
- There are no good quality studies to determine the efficacy and safety of hydroxyurea
- In a prospective observational study, hydroxyurea 500 mg twice daily was an efficacious **alternative** treatment for patients with chronic plaque psoriasis
 - 76% of patients had a PASI 75 reduction at week 10 - 12

Hydroxyurea

- Adverse events reported were
 - Leucopenia
 - Thrombocytopenia
 - Skin infection
 - Dry skin
 - Diffuse reversible alopecia
 - Anaemia
 - Post-inflammatory lesional and
 - Nail hyperpigmentation were seen in all patients taking hydroxyurea

Salazopyrin (Sulfasalazine)

- Salazopyrin is a 5-lipoxygenase inhibitor which has anti-inflammatory and immunomodulatory effect
- It had been shown to be efficacious in treating psoriatic arthritis based on several randomized controlled trials in 1990s
- Evidence on the use of salazopyrin in treating plaque psoriasis is limited.

Salazopyrin (Sulfasalazine)

- 8 week double-blind randomized controlled trial, salazopyrin (1.5 g to 4.0 g)(n=23) vs placebo(n=27) was more efficacious to achieve improvement
 - marked improvement : 41% vs 0%
 - moderate improvement : 41% vs 4%
- In the salazopyrin arm
 - 26% dropped out at the end of 8 week due to rash or nausea
 - 14 patients who continued for an additional 4 weeks
 - 8 had marked improvement
 - 2 had moderate improvement

Leflunomide

- Leflunomide is a dihydro-orotate dehydrogenase inhibitor which is a key enzyme in the de novo synthesis of pyrimidine
- Limited randomized controlled trial on the efficacy of leflunomide
- In a randomized double blind placebo controlled trial
 - leflunomide 20mg (n=95) vs placebo (n=91)
 - Leflunomide was more efficacious
 - PASI 75 : 17.4% vs 7.8% (p=0.048)

Leflunomide

- In the same study, leflunomide showed higher incidences of
 - diarrhea (24%)
 - increased alanine transaminase level (12.5%)
 - lethargy (6.3%)

As compared to placebo

Malaysian CPG on the management of Psoriasis vulgaris

Systemic therapy

RECOMMENDATION

- All patients for systemic/biologic therapy should have a pre-treatment assessment including laboratory/imaging tests and regular monitoring for side-effects/toxicity. **(Grade C)**
- Methotrexate should be used as first-line systemic treatment for moderate to severe plaque psoriasis. **(Grade A)**
 - Neutropaenia and hepatotoxicity should be monitored closely. **(Grade C)**
- Acitretin may be offered for the treatment of moderate to severe plaque psoriasis. **(Grade A)**
 - Acitretin should be avoided in women of childbearing age without reliable contraception and in those who are planning pregnancy, however, it is safe for men who are planning to have a child. **(Grade C)**

**Malaysian CPG on the management of
Psoriasis vulgaris
Systemic therapy**

RECOMMENDATION

- Cyclosporine may be offered as short-term treatment for rapid disease clearance in moderate to severe psoriasis. **(Grade A)**
- Cyclosporine may be offered as second-line systemic agent to psoriasis patient who fail, intolerant or have **contraindications to methotrexate. (Grade A)**
 - Cyclosporine should **NOT** be used for more than 2 years. **(Grade B)**
 - Cyclosporine should be avoided in patient with previous PUVA exposure. **(Grade B)**
 - Blood pressure, renal function, lipid profile should be monitored closely in psoriasis patient on cyclosporine. **(Grade C)**