SUMMARY OF MALAYSIAN NATIONAL CANCER REGISTRY REPORT 2007-2011

THIS BOOKLET PROVIDES SUMMARIZED INFORMATION ON CANCER FACTS IN MALAYSIA BASED ON THE FIRST 5 YEARS REPORT (2007-2011) OF THE NATIONAL CANCER REGISTRY (MNCR)

SUMMARY OF MALAYSIAN NATIONAL CANCER REGISTRY REPORT 2007-2011







NATIONAL CANCER REGISTRY DEPARTMENT NATIONAL CANCER INSTITUTE, MINISTRY OF HEALTH

No. 4, Jalan P7, Precint 7, 62250 Putrajaya Tel: (603)88925555 Fax: (603)88925625

DIRECTOR OF NATIONAL CANCER INSTITUTE

Dr. Asmayani Binti Khalib

STAFF MEMBERS OF THE REGISTRY:

HEAD OF NATIONAL CANCER REGISTRY DEPARTMENT

Datin Dr. Azizah Binti Ab Manan, Public Health Physician Email: drazizah@nci.gov.my

MEDICAL OFFICERS

Dr. Puteri Nureylia Binti Amir Email: drnureylia@nci.gov.my Dr. Renuga A/P K. Raman Email: drrenuga@nci.gov.my


Dr. Nabihah Binti Ali Email: drnabihah@nci.gov.my Dr. Sukumaran A/L Raman Email: drsukumaran@nci.gov.my

ASSISTANT ADMINISTRATOR

Puan Nor Hafizah Binti Mohd Nor Email: ptnorhafizah@nci.gov.my

TEN MOST COMMON CANCER IN MALAYSIA, 2007-2011

(PERCENTAGES)




TEN MOST COMMON CANCERS IN MALAYSIA BY SEX, 2007-2011

MALE				FEMALE					
ICD 10	SITE	NO.	%	ICD 10	SITE	NO.	%		
C18-C21	Colorectal	7,646	16.4	C50	Breast	18,206	32.1		
C33-C34	Lung, Trachea, Bronchus	7,415	15.8	C18-C21	Colorectal	6,047	10.7		
C11	Nasopharynx	3,785	8.1	C53	Cervix Uteri	4,352	7.7		
C81-C85, C96	Lymphoma	3,171	6.8	C56	Ovary	3,472	6.1		
C61	Prostate	3,132	6.7	C33-C34	Lung, Trachea, Bronchus	3,193	5.6		
C22	Liver	3,054	6.5	C81-C85, C96	Lymphoma	2,203	3.9		
C91-C95	Leukaemia	2,549	5.4	C54	Corpus Uteri	2,181	3.8		
C16	Stomach	2,014	4.3	C91-C95	Leukaemia	2,024	3.5		
C67	Bladder	1,477	3.2	C73	Thyroid	1,723	3.0		
C44	Other Skin	1,384	3.0	C16	Stomach	1,447	2.6		



INCIDENCE RATE (ASR) FOR TEN MOST COMMON CANCERS BY SEX


AGE STANDARDISED RATE (WORLD) PER 100.000 POPULATION



Definition of Age-Standardised Rate (ASR): The age-standardised incidence rate is a summary measure, indicating the rate that a population would have if it had a standard age structure. It is calculated by summing the age-specific rates weighting to the world standard population; the calculated incidence rate is then called the World Standardised incidence rate. It is also expressed per 100,000.

LIFETIME RISK


THE PROBABILITY THAT AN INDIVIDUAL WILL DEVELOP THE CANCER BEFORE THE AGE OF 75 IN THE ABSENCE OF OTHER CAUSES OF DEATH.






LIFETIME RISK

MAJOR ETHNIC GROUPS





AGE-SPECIFIC INCIDENCE RATE FOR ALL CANCERS



The incidence of cancer increased for both male and female after the age of 30 years. The incidence rate in males exceeded the rate in females after the age of 60 years.

Definition of Age-specific Rate (AR): Incidence rate in a specified five-year age group; calculated by dividing the number of cancer cases in the five-year age group by the Malaysian population in the particular age group and multiply by 100,000.



9 SELECTED CANCERS





Breast

Colorectal

Lung, Trachea, Bronchus

Nasopharynx

Cervix Uteri

Ovary



Prostate

Stomach


Corpus Uteri






FEMALE BREAST CANCER

AGE SPECIFIC INCIDENCE RATE



Breast cancer was the **most common** cancer in Malaysia. The incidence was **highest among Chinese** followed by Indian and Malay. Most of the cases were presented at the age of 45-69 years and reduced after the age of 70 years.


THE LIFETIME RISK: 1 in 30 for all females | 1 in 35 Malay | 1 in 22 Chinese | 1 in 24 Indian



Staging was reported for 11,938 cases (66%) of these, 43% was diagnosed at very late stage (Stage 3 & 4)

COLORECTAL CANCER


AGE SPECIFIC INCIDENCE RATE



Colorectal cancer was the **most common** cancer in **male** and second in female. The incidence increased after the age of 60 years in both sexes

THE LIFETIME RISK:


1 in 56 for all male | 1 in 72 for Malay | 1 in 39 for Chinese | 1 in 82 for Indian 1 in 74 for all female | 1 in 100 for Malay | 1 in 49 for Chinese | 1 in 91 for Indian




(Stage 3 & 4) for both sexes (66% in male; 65% in female)

LUNG, TRACHEA & BRONCHUS CANCER

AGE SPECIFIC INCIDENCE RATE



These cancers were the **second most common** cancer among **male** and fifth among female. The incidence was higher in male and started to peak at the age of 60 years old.



Extremely high percentages of the lung, trachea, bronchus cancer cases were detected at late stage (Stage 3 & 4) for both sexes (89% in male; 91% in female)

STAGE 4 | 71%

SELECTED CANCERS

6

STAGE 4 | 66%


NASOPHARYNX CANCER

AGE SPECIFIC INCIDENCE RATE






Nasopharyngeal cancer was the fifth most common cancer and **third most common** in male. The incidence was higher in male and the trend started to increased from the age of **45** years and **highest** at the age of **65-69** years old.




(Stage 3 & 4) for both sexes (63% in male; 60% in female)

CERVIX UTERI CANCER

AGE SPECIFIC INCIDENCE RATE




Cervix uteri cancer was the **third most common cancer in female**. The incidence trend increased at the age of 35 years and **peak between 50-74** years old.




SELECTED CANCERS

OVARIAN CANCER

AGE SPECIFIC INCIDENCE RATE




Ovarian cancer was the **fourth common cancer in female**. The incidence trend increased at the age of 35 years and **peak between 45-69 years old** and **reducing trend after 70** years.



SELECTED CANCERS


STOMACH CANCER

AGE SPECIFIC INCIDENCE RATE





Stomach cancer was eight most common cancer in male and tenth in female. The incidence increased with age and **peak above 60 years old** for both sexes.



stage (Stage 3 & 4) for both sexes (75% in male; 74% in female).


PROSTATE CANCER

AGE SPECIFIC INCIDENCE RATE

AGE 0-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50-54 55-59 60-64 65-69 70-74 75+


MALE

Prostate cancer was the fifth common cancer in male. The incidence trend increased at the age of 55 years and **mostly diagnosed after the age of 65 years old**.




CORPUS UTERI CANCER

AGE SPECIFIC INCIDENCE RATE



Corpus uteri cancer was the seventh most common cancer in female. The incidence trend increased at the age of 40 years and the **highest incidence** was noted between the age of **55-64 years** and reduced after the age of 65 years old.



INCIDENCE SUMMARY TABLES

Incidence Sum	mary by Sex,	All residents	2007-2011
---------------	--------------	---------------	-----------

Sex	Number	%	CR	ASR	CumR
MALE	46,794	45.2	71.4	86.9	9.8
FEMALE	56,713	54.8	89.0	99.3	10.6
Both	103,507	100	1111		

	Incidence Summary by Ethnic Groups and Sex, 2007-2011							
	MALE				FEMALE			
Ethnicity	Number	CR	ASR	CumR	Number	CR	ASR	CumR
Malay	18,757	53.1	70.7	8.0	24,188	69.5	81.9	8.7
Chinese	20,424	124.4	112.9	12.5	22,114	143.2	126.2	13.2
Indian	2,679	56.0	70.4	8.1	4,798	101.2	110.1	12.2
Others*	4,934	55.0	80.5	9.2	5,613	64.4	89.6	9.7

	Incidence Summary by Year and Sex								
	MALE					FEMALE			
Year	Number	CR	ASR	CumR	Number	CR	ASR	CumR	
2007	8,969	70.8	89.7	10.1	11,035	89.5	102.6	10.8	
2008	9,383	72.5	90.2	10.1	11,344	89.7	102.3	10.9	
2009	9,495	72.5	87.9	9.9	11,216	88	98.4	10.6	
2010	9,571	71.9	85.4	9.7	11,672	90.1	98.8	10.6	
2011	9,376	68.6	80.9	9.3	11,446	87.2	93.8	10	

Incidence Summary by State and Sex, 2007-2011

		MALE		FEMALE			
State	Number	CR	ASR	Number	CR	ASR	
Johor	7,202	91.8	109	8,110	109.4	118.7	
Selangor	6,261	49.3	71.3	8,484	69.2	89.7	
Penang	5,245	144.1	150.5	6,085	166.6	157.8	
Sarawak	4,549	76.4	85.7	5,185	90.3	100.2	
Perak	4,465	77.9	74.7	5,504	96.6	88.4	
Sabah	3,522	58.9	93.6	4,045	69.8	105.3	
Kedah	2,774	59.6	66.5	3,565	76.7	77	
K. Lumpur & Putrajaya	2,277	59.5	73.1	3,307	88.3	101.2	
Pahang	2,193	60.9	75.1	2,499	73.6	82.6	
Kelantan	2,105	54.8	68.0	2,281	59.6	65.8	
Negeri Sembilan	1,917	79	89.1	2,528	106.9	108.9	
Melaka	1,811	94.2	104.7	2,214	110.2	108.1	
Terengganu	1,668	63.7	83.1	1,977	79.1	93.8	
Perlis	405	46.2	72.8	495	86.4	84.9	
Labuan	53	26.5	50.9	51	26.8	41.7	

Definition of Crude incidence Rate (CR): The number of new cancer cases (incidence cases) observed in the population during a defined period, divided by the number of population at risk in the same period. It is usually expressed per 100,000.

Definition of Cumulative Risk (CumR): Cumulative risk is defined as a probability that an individual would develop cancer during a certain age period, in the absence of any competing cause of death. The age period over which the risk is accumulated in this report is 0-74 years.



MINISTRY OF HEALTH

Published by National Cancer Registry Department Collaboration with Health Education Unit National Cancer Institute Ministry of Health Putrajaya http://nci.moh.gov.my ncr.putrajaya@nci.gov.my

3000/01/2017